

# Stratamed – the breakthrough treatment for the use on open wounds and compromised skin

**Stratamed can be used immediately after surgical procedures, trauma, skin resurfacing, burns\* and radiation therapy.**

- Allows prevention of abnormal scarring to begin earlier than ever before
- Faster wound healing<sup>1</sup>
- Faster recovery time

**Stratamed creates an optimal environment for the healing process of the skin, which leads to:**

- Dramatically decreased incidence of abnormal scarring<sup>2</sup>
- Reduction in the inflammatory response<sup>1</sup>
- Reduced symptomatic responses<sup>1</sup> (redness/discoloration, itching, discomfort and pain)
- Enhanced therapeutic results<sup>1</sup>

**Visit our website for more information:**  
[www.stratamed.com](http://www.stratamed.com)

**Simple, safe and convenient**

- Easy to use at home
- Fast drying
- Can be used on hairy areas and areas with contours
- Can be covered with make-up or sunscreen

Stratamed helps you to improve the visible outcome after skin damage, prevents abnormal scarring and reduces the healing time of your wound!



Strataderm is available in 5g, 10g, 20g and 50g tubes.

\* First and second degree burns, third degree burns only after the presence of granulating tissue

1. Data on file. Stratapharma AG; Practice Dr. Sandhofer, Linz/Austria 2010; 2. Gauglitz GG et al. Mol Med 2011;17(1-2):113-125

**Fresh wound post incision<sup>1</sup>** Monotherapy treatment with Stratamed.



Before treatment



Week 8

**Laser resurfacing (wrinkles)<sup>1</sup>** Treatment with Stratamed in combination with thermage, CO<sub>2</sub> fractional laser resurfacing and Botox.



Before treatment



Day 0



Day 10

**Dermabrasion<sup>1</sup>** Monotherapy treatment with Stratamed.



During treatment



During treatment



Week 4